

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
Please do not report the images to the
Image Problem Mailbox.

(12) UK Patent Application (19) GB (11) 2 018 593 - A

- (21) Application No 7914799
(22) Date of filing
27 Apr 1979
(23) Claims filed
27 Apr 1979
(30) Priority data
(31) 3858/78
(32) 31 Jan 1975
(33) United Kingdom (GB)
(43) Application published
24 Oct 1979
(51) INT CL² A01N 17/14
(52) Domestic classification
A5E 405 409 410 411
412 503 510 K
(56) Documents cited
GB 1480943
GB 1440860
GB 1391248
GB 1182007
GB 1145433
GB 955680
GB 987173
(59) Field of search
A5E
(71) Applicant
Unilever Limited
Unilever House
Blackfriars
London EC4
England
(72) Inventor
Norman Albert Hunt
(74) Agents
Mr J E Farndon


(54) Insect control systems


(57) This invention relates to insect control systems comprising insect attractants and insecticides in carrier compositions formulated to ensure comparable effective lives for the two compositions. The carriers may be P.V.C., P.V.A. or acetate chloride copolymers, polyolefins, chlorinated polyethylene, urea and melamine formaldehyde resins, polyesters, polyurethanes, polyureas, gelatins, straw, cane, lignocellulose, silica, aluminosilicates or clays, or the active substances may be microencapsulated. The compositions may be in the form of tapes or strings, or particles disposed on a polymeric sheet. The system may be used in conjunction with an insect trap.

GB 2 018 593 A

2018593


GRAPH 1-EXAMPLE LIFE AND RELEASE OF ATTRACTANT AND INSECTICIDE
IN BALANCED CONTROLLED RELEASE "ATTRACT & KILL" PROCEDURES





2018593

GRAPH 2 - EXAMPLE II. LIFE AND RELEASE OF ATTRACTANT AND INSECTICIDE
IN BALANCED CONTROLLED RELEASE "ATTRACT & KILL" PROCEDURES



SPECIFICATION

Insect control systems

- 5 This invention relates to insect control systems. More particularly, the invention relates to
an insect control system which optimises and generally minimises the use of insecticides
and can, in preferred forms, be species specific with regard to the insects affected by the
system. 5
- The term "insecticide" as used in this specification is intended to include orthodox
10 chemical insecticides and appropriate insect virus, bacterial or hormone compositions able
to affect the specific insect species under attack. The insecticide can be of the contact type,
or one which is effective in the form of its vapour. 10
- Insecticides have been used for many years to combat various insect species which cause
damage to crops. Insecticides are frequently dangerous and persistent chemicals and,
15 hitherto, they have usually been applied to crops by dusting or spraying, either from the
ground or from the air, the insecticide being directed to the space occupied by the crop to
be protected and its surroundings. Hence, larger quantities of insecticides are used than
would be used if the insecticide could be applied directly to, and only to, the target insect. 15
- In addition to the large quantities of insecticides used, such application techniques are
20 indiscriminate in that all insect species present and the crop itself are contaminated with
significant quantities of insecticide. Insecticides so broadcast will affect both useful insects,
such as pollinators and also insect predators, which attack the harmful or target insect and
can, in many circumstances, be counter-productive. 20
- Since the insecticide will directly contact the crop which is to be protected, if the crop is
25 a food crop, it is normally not safe to apply the insecticide for many days immediately prior
to harvesting the crop. 25
- Finally, many insecticides are persistent and applying large quantities of such insecticides
to areas of land can create a long-term pollution problem, the consequences of which are
not fully understood.
- 30 It is known that in many insect species behaviour of the insect is influenced by certain
specific volatile substances and mixtures of substances. Such chemical substances may, for
example, be emitted by the female insect and serve to indicate her location to the male,
which travels to the source of the substance. In other cases, volatile substances from
appropriate host plants will direct the female to lay her eggs on such plants. In addition, it
35 has been found that certain volatile substances are attractive to particular insect species,
although the exact significance of the attraction is not fully understood. In many of these
cases the volatile substance is species specific—that is to say, attracts one species of
insect. These volatile substances are sometimes known as pheromones, lures or attractants
and can attract insects over significant distances. All these substances are hereinafter
40 referred to as insect attractants. 40
- It has now been found that insecticides and insect attractants can be incorporated into
compositions from which they can be discharged at a predetermined rate and the present
invention provides an insecticidal system which avoids the need to scatter large quantities
of insecticide over growing crops and can be used to attract specific insects to an
45 appropriate insecticide, hence reducing the risk of damaging non-harmful insects. 45
- Accordingly, the present invention provides an insect control system comprising an
assembly of at least one insecticidal carrier composition and one insect attractant carrier
composition, said compositions being formulated to provide effective lives for the composi-
tions of substantially the same period.
- 50 In a preferred form of the invention, the effective life of the insecticide is longer than that
of the insect attractant so that under no circumstances will the attractant exist in the
absence of the insecticide. 50
- The insecticide and insect attractant incorporated into carrier compositions can be
mounted in an insect trap or juxtaposed compositions can be prepared: one containing the
55 insecticide and the other the attractant, for example as intertwined strings or tapes. 55
- The carrier composition can conveniently be a polymer plastics material which will
release the attractant and insecticide over a period of some weeks, or alternatively, if a fairly
fast release rate is required, a cellulosic material, such as compressed paperboard, may be
employed. Conveniently, the carrier composition is biodegradable, although if the insectici-
dal system is to be employed in association with an insect trap or container, then it is
60 frequently convenient to use a polymeric material which can be mounted in an appropriate
container or trap. In addition a silica gel adsorbate may be used as the carrier for either or
both of the compositions. 60
- Conveniently, the carrier composition comprises a polymeric sheet material incorporating
65 the insecticide and carrying, on at least one surface, a secondary carrier composition 65

TABLE I

5	Attractant (Lure)	Chief Insect Pests Against which Lure is Used	5
10	1. Cue-lure* 4-(p-acetoxy phenyl)butan-2-one	Dacus cucurbitae (Coquillett) (Melon Fruit Fly) Dacus tryoni (Froggatt) (Queensland Fruit Fly) Dacus dorsalis (Hendel) (Oriental Fruit Fly) Ceratitis capitata (Wiedemann) (Mediterranean Fruit Fly) Trichoplusia (Hubner) (Cabbage Looper) Laspeyresia pomonella (Linnaeus) (Codling Moth) Adoxophyes orana (Summer Fruit Tortrix Moth)	10
15	2. Methyl eugenol 4,5 dimethoxy propenyl benzene		15
20	3. Trimedlure Isobutyl ester of 2 methyl (4/5) chloro cyclohexane carboxylic acid		20
25	4. (Z)-7-dodecan-1-yl acetate		25
30	5. (E,E)-8-10 dodecadien-1-ol		30
35	6. (Z)-9-tetradecenyl acetate (E)-9-tetradecenyl acetate (Z)-11-tetradecenyl acetate (E)-11-tetradecenyl acetate		35
40	7. (E)-11-tetradecenal	Choristoneura fumiferana (Clemens) (Spruce Budworm) Choristoneura occidentalis (Freeman) (Western Spruce Budworm) Spodoptera exempta (Walk) (Army Worm)	40
45	8. Z-9-tetradecenyl acetate 20 parts (Z)-9, (E)-12 tetradecadien-1-yl acetate 1 part		45
50	9. (Z,E)-9,11-tetradecadien-1-yl acetate	Spodoptera littoralis (Boisd) (Egyptian Cotton Leaf Worm) Heliothis virescens (Fabricius) (Tobacco Budworm) Ceratitis capitata (Wiedemann) (Mediterranean Fruit Fly) Spodoptera frugiperda (J.E. Smith) (Fall Armyworm Moth) Prodenia eridania (Cramer) (Southern Armyworm Moth) Pectinophora gossypiella (Saunders) (Pink Bollworm) Heliothis zea (Boddie) (Bollworm, Corn Ear Worm, Tomato Fruit Worm) Sesamia inferens (Walk) (Purple Stem Borer Moth) Chilo suppressalis (Walk) (Striped Rice Borer) Porthetria diapa (Lo) (Gypsy Moth)	50
55	10. (Z)-9-tetradecenal (Z)-11-hexadecenal		55
60	11. Capilure*		60
	12. Z-9-tetradecen-1-ol acetate (Z,E)-9,12-tetradecadien-1-ol acetate		
	13. (Z)-7, hexadecen-1-yl acetate		
	14. (Z)-11 hexadecenal		
	15. (Z)-11-hexadecenyl acetate		
	16. Z-11-hexadecenal-5-parts (Z)-13-octadecenal 1 part		
	17. Disparlure cis 7, 8 poxy-2-methyl- octadecane		

(*Trade Mark of Food Industries Limited Bromborough, England.)

invention comprising a PVC-based carrier composition. An attractant carrier composition was prepared using:

	Parts by weight	
5		5
	Breon, P 130/1 (PVC Emulsion Polymer -BP Limited)	100.0
	DOP (Diethyl Phthalate-Plasticiser)	54.0
10	Vimco 249 C (Barium/Cadmium-Stabiliser)	2.5
	ED6 (Epoxy Stabiliser-Lankro Chem. Co.)	5.0
	Tinuvin P (UV Adsorber-Ciba-Geigy)	0.1
	Pigment (Phthalocyanine Green or Chrome Yellow-according to attractant)	0.5
15	Insect Attractant	3.3
		15

Approximately 15% of the plasticiser was charged into a mixer and the PVC polymer emulsion then added, followed by the remainder of the constituents in the order listed above. The balance of the plasticiser was then added, followed by the remainder of the constituents in the order listed above. The balance of the plasticiser was then added and mechanical mixing commenced, first at low speed and then at a higher speed, until a smooth paste was obtained which was passed through a triple-roll mill. The resultant paste was spread to a desired thickness (approximately 2 mm) onto a release paper and heated to 180 to 200°C for 3 minutes and then cooled for 10 minutes. This carrier composition contained approximately 2% by weight of the insect attractant.

Using a similar technique, an insecticide-containing carrier composition was prepared from the following formulation:

	Parts by weight	
30		30
	Vinmol E10/65F (PVC Emulsion Polymer)	100.0
	B.Br. (Butyl Benzyl Phthalate Plasticiser)	60.0
35	Vimco 249 C (Barium/Cadmium Stabiliser)	2.5
	ED6 (Epoxy Stabiliser-Lankro Chem. Co.)	5.0
		35

	Parts by weight	
40		40
	Tinuvin P (UV Adsorber-Ciba-Geigy)	0.1
	Pigment (Azo Red)	0.5
	Insecticide	42.0
45		45

This yielded a carrier composition containing approximately 20% by weight of insecticide.

50 Example 1

A carrier composition comprising, as an insect attractant, methyl eugenol, which is an attractant for the Oriental Fruit Fly, was prepared as described above. The carrier composition contained 2% by weight of the methyl eugenol.

A further carrier composition comprising insecticide was prepared, this time using, as insecticide, a mixture of equal parts by weight of:

	Dichlorvos	2,2-dichlorovinyl-dimethyl phosphate	
	Naled. Dibrom	1,2-dibromo-2,2-dichloro ethyl dimethyl phosphate	
60	Malathion	0,0-dimethyl S-diethyl-mercapto succinate phosphorodithioate	60

This insecticidal carrier composition contained 20% by weight of composition of the mixture of insecticides.

Constituents	Composition % by weight				
5 Polyvinyl chloride PVC Corvic D65/02	55.33	5			
DIDP Plasticiser (Drisodecylphthalate)	29.57				
Mark 33 (Calcium/zinc oxides) Stabilizer	1.38				
Mark C Antioxidant (Trinonyl phenyl phosphite)	0.28				
10 Paraplex G62 (Stabiliser/plasticiser) Epoxy		10			
Soya Bean Oil	2.77				
Calcium Stearate (Lubricant)	0.18				
Ultra-violet Adsorber Tinuvin P	0.06				
Red Pigment (Vinamon G)	0.28				
15 Insecticide (Mixture of equal parts of Dichlorvos, Dibrom and Malathion)	10.15	15			
	100.00				
20		20			
<i>Example V</i>					
<i>Controlled Release—Lure Compositions</i>					
An alternative method of obtaining the desired controlled release rates of the lures and the insecticides, is illustrated:					
25	The selected lure (Cue-lure, methyl eugenol or Trimedlure) was cold mixed into a matrix of an adhesive polymer composition based on acrylic polymer blends 50% and iso propyl acetate 50% (exemplified by Adhesive 5050 of Vinyl Products Limited, Carshalton, Surrey England). The intimately mixed adhesive and lure were spread onto 2 metre unit lengths of one of the following polymer films—each as a separate formulation.	25			
30	Polythene Polyvinyl chloride Terephthalate polyester	30			
35	These adhesive covered polymer films were then covered with a further polymer film of the same composition. These completed plastic sandwiches were approximately of dimensions 200 cm × 5 cm × 0.3 cm. Adjuncts such as the antioxidants, ultra-violet screen compounds and dyes were included as required. <i>Lure contents</i> (Cue-lure, methyl eugenol or Trimedlure) per unit 2 metre length	35			
40	plastic sandwich strip were: 13.2 to 13.8 grammes	40			
<i>Controlled Release—Insecticide Compositions</i>					
45	In this example, a layer of insecticide, adhesive polymer mixture of composition:	45			
		% by weight			
50	Dichlorvos	37.5	50		
	Polymer (5050)	62.5			
was spread evenly between two plastic strips (Polyester Terephthalate) Polymer/Insecticide layer was 5 cm wide.					
55	Insecticide composition was:			55	
60	Concentration	Per cubic centimetre.	Per square centimetre	Per Centimetre length	60
	<i>Insecticide</i> Dichlorvos	200.0 mg	8.0 mg	40.0 mg	

The weight composition of the system is:

	Grammes	Percent	
5			5
Total weight of a test strip	6.5	100.0	
Weight of plastic film	3.0	46.2	
Weight of adhesive polymer (5050)	2.2	33.8	
Weight of insecticide	1.3	20.0	
10			10

The properties of the film compositions used as illustrated by their vapour permeability properties for water vapour and oxygen.

	Thickness		Permeability Value				
	ins	mm	Test Temp. °C	Test Relative Humidity %	Water Vapour gms/m ²	Oxygen cc/m ²	
20							20
Composition							
Polythene	0.001	0.025	25	—	1.0	350.0	
PVC	—	0.013	25	75	2.0	192 × 106*	
PVC	0.02	0.51	—	—	—	—	
25							25
Terephthalate (Polyester)	0.001	0.025	38	90	0.90	19.0	

*Test carried out at 21°C and 44% relative humidity.

- 30 In the accompanying drawings:
 Graphs I and IA show the release of the active components of Example I.
 Graph 2 gives similar information in relation to Example II.

CLAIMS

- 35 1. An insect control system comprising an assembly of at least one insecticidal carrier composition and one insect attractant carrier composition, said compositions being formulated to provide effective lives for the compositions of substantially the same period.
2. An insect control system as claimed in Claim 1 in which the effective life of the insecticidal composition is longer than that of the attractant composition.
- 40 3. An insect control system as claimed in Claim 1 or 2, in which the assembly of compositions is juxtaposed.
4. An insect control system as claimed in Claim 3 in which the juxtaposed compositions are in the form of interwoven tapes or strings.
5. An insect control system as claimed in Claim 3 in which one carrier composition
- 45 comprises a polymeric sheet material incorporating the insecticide and carrying, on at least one surface, a secondary carrier composition containing the insect attractant, the secondary composition being in the form of discrete particles.
6. An insect control system as claimed in Claim 5 in which the discrete particles formed from the secondary carrier composition and the insect attractant are in the range of
- 50 diameters up to half the length of the insect under attack.
7. An insect control system as claimed in Claim 3 in which one of the compositions is in micro-encapsulated form and carried on the surface of the other composition.
8. An insect control system as claimed in any one of the preceding claims in which the assembly is located in an insect trap.
- 55 9. An insect control system, substantially as described herein, with reference to the Examples.